Turkmenistan


2010 total population: 5 041 995 Income group: Lower middle

NCD mortality		
2008 estimates	males	females
Total NCD deaths (000s)	16.1	15.7
NCD deaths under age 60 (percent of all NCD deaths)	50.4	32.0
Age-standardized death rate per 100 000		
All NCDs	1181.7	872.8
Cancers	120.6	92.1
Chronic respiratory diseases	49.3	32.0
Cardiovascular diseases and diabetes	8.088	667.7

Behavioural risk factors			
2008 estimated prevalence (%)	males	females	total
Current daily tobacco smoking			
Physical inactivity			


Metabolic risk factors			
2008 estimated prevalence (%)	males	females	total
Raised blood pressure			
Raised blood glucose			
Overweight	44.8	38.5	41.5
Obesity	12.9	13.5	13.2
Raised cholesterol			

Proportional mortality (% of total deaths, all ages)



NCDs are estimated to account for 73% of all deaths.

Metabolic risk factor trends



Country capacity to address and respond to NCDs			
Has a Unit / Branch / Dept in MOH with responsibility for NCDs	ND	Has an integrated or topic-specific policy / programme plan which is currently operational for:	e / action
There is funding available for:		Cardiovascular diseases	ND
NCD treatment and control	ND	Cancer	ND
NCD prevention and health promotion	ND	Chronic respiratory diseases	ND
NCD surveillance, monitoring and evaluation	ND	Diabetes	ND
		Alcohol	ND
National health reporting system includes:		Unhealthy diet / Overweight / Obesity	ND
NCD cause-specific mortality	ND	Physical inactivity	ND
NCD morbidity	ND	Tobacco	ND
NCD risk factors	ND		
Has a national, population-based cancer registry	ND	Number of tobacco (m)POWER measures implemented at the highest level of achievement	1/5

^{.. =} no data available

ND = Country did not respond to country capacity survey